

Some Unsettled Questions in the Problem of Neutrino Oscillations. Remarks About the Majorana Neutrino Oscillations

Beshtoev Kh. M.

Joint Inst. for Nucl. Research., Joliot Curie 6, 141980 Dubna, Moscow region, Russia; Inst. of Appl. Math. and Autom. of the Kabardino-Balkarian Scientific Center of RAC, Shortanova 89a, 360017 Nalchik, Russia

Abstract

It is shown that we cannot put Majorana neutrinos in the standard Dirac theory without violation of the gauge invariance. Experiments on $\nu_l \rightarrow l (l = e, \mu, \tau)$ transitions do not confirm this supposition. It means that on neutrino oscillation experiments, the Majorana neutrino oscillations cannot be observed.

1. Introduction

At present, it is supposed [1, 2] that the neutrino oscillations can be connected to Majorana neutrino oscillations. This idea is based on suggestion that the neutrinos are the Majorana ones i.e. that the standard weak interactions theory includes only these neutrinos. Let us come to proof that it leads to contradictions in the theory and is not confirmed on experiments.

2. Majorana Neutrino Oscillations

Majorana fermion in Dirac representation has the following form [1, 3, 4]:

$$\begin{aligned}\chi^M &= \frac{1}{2}[\Psi(x) + \eta_C \Psi^C(x)], \\ \Psi^C(x) &\rightarrow \eta_C C \bar{\Psi}^T(x),\end{aligned}\tag{1}$$

where η_C is a phase, C is a charge conjunction, T is a transposition.

From Exp. (1) we see that Majorana fermion χ^M has two spin projections $\pm\frac{1}{2}$; and then the Majorana spinor can be rewritten in the following form:

$$\chi^M(x) = \begin{pmatrix} \chi_{+\frac{1}{2}}(x) \\ \chi_{-\frac{1}{2}}(x) \end{pmatrix}. \quad (2)$$

The mass Lagrangian of Majorana neutrinos in the case of two neutrinos χ_e, χ_μ ($-\frac{1}{2}$ components of Majorana neutrinos and $\bar{\chi}_{\dots}$ are the same Majorana fermion with the opposite spin projection) in the common case has the following form:

$$\mathcal{L}'_M = -\frac{1}{2}(\bar{\chi}_e, \bar{\chi}_\mu) \begin{pmatrix} m_{\chi_e} & m_{\chi_e\chi_\mu} \\ m_{\chi_\mu\chi_e} & m_{\chi_\mu} \end{pmatrix} \begin{pmatrix} \chi_e \\ \chi_\mu \end{pmatrix}. \quad (3)$$

By diagonalizing this mass matrix by standard methods, one obtains the following expression:

$$\mathcal{L}'_M = -\frac{1}{2}(\bar{\nu}_1, \bar{\nu}_2) \begin{pmatrix} m_{\nu_1} & 0 \\ 0 & m_{\nu_2} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}, \quad \begin{aligned} \nu_1 &= \cos\theta\chi_e - \sin\theta\chi_\mu \\ \nu_2 &= \sin\theta\chi_e + \cos\theta\chi_\mu \end{aligned}. \quad (4)$$

These neutrino oscillations are described by standard expressions (see [1,4,5]).

The standard theory of weak interactions is constructed on the base of local gauge invariance of Dirac fermions. In this case, Dirac fermions have the following lepton numbers l_l , which are conserved, $l_l, l = e, \mu, \tau$, and Dirac antiparticles have lepton numbers with the opposite sign $\bar{l} = -l_l$.

Gauge transformation of Majorana fermions can be written in the following form:

$$\chi'_{+\frac{1}{2}}(x) = \exp(-i\beta)\chi_{+\frac{1}{2}}(x), \quad \chi'_{-\frac{1}{2}}(x) = \exp(+i\beta)\chi_{-\frac{1}{2}}(x). \quad (5)$$

Then, lepton numbers of Majorana fermions are

$$l^M = \sum_i l_i^M(+1/2) = -\sum_i l_i^M(-1/2), \quad (6)$$

i. e., antiparticle of Majorana fermion is the same fermion with the opposite spin projection.

Now, we come to discussion of the problem of the place of Majorana fermion in the standard theory of weak interactions [6].

In order to construct the standard theory of weak interactions [7], Dirac fermions are used. In this theory, the fermions are present as doublets. The local current $j^{\mu i}$ of the weak interaction has the following form:

$$j^{\mu i} = \bar{\Psi}_L \tau^i \gamma^\mu \Psi_L, \quad (7)$$

where $\bar{\Psi}_L, \Psi_L$ are lepton doublets

$$\begin{pmatrix} e \\ \nu_e \end{pmatrix}_{iL} \quad i = 1 - 3, \quad (8)$$

where i is flavor number of leptons. If we consider the process transition of neutrino to corresponding lepton, we will get the following reaction:

$$\nu_l + A(Z) \rightarrow l^- + A(Z + 1) \quad (9)$$

The absence of contradictions in this theory with the experimental data (see [8]) confirms that all fermions are Dirac particles.

Now, if we want to put the Majorana fermions into the standard theory, we must take into account that, in the common case, the gauge charges of the Dirac and Majorana fermions are different (especially well it is seen in the example of Dirac fermion having an electrical charge since it cannot have a Majorana charge). In this case we cannot just include Majorana fermions in the standard theory of weak interactions by gauge invariance manner. Then, in the standard theory the Majorana fermions cannot appear.

In spite of above arguments, if we put Majorana neutrinos in the standard theory; then on experiments we must see the following reactions:

$$\chi_l + A(Z) \rightarrow l^- + A(Z + 1)$$

with relative probability 1/2 and

$$\chi_l + A(Z) \rightarrow l^+ + A(Z - 1)$$

with the same relative probability (where $x = e, \mu, \tau$), since Majorana neutrinos are superpositions of Dirac neutrinos and antineutrinos. Obviously, all the available experimental data [8] do not confirm this predictions; therefore we cannot consider this mechanism as a realistic one for neutrino oscillations.

3. Conclusion

It was shown that we cannot put Majorana neutrinos in the standard Dirac theory without violation of the gauge invariance. Experiments on $\nu_l \rightarrow l (l = e, \mu, \tau)$ transitions do not confirm this supposition. It means that on neutrino oscillation experiments the Majorana neutrino oscillations cannot be observed.

References

1. Bilenky S.M., Pontecorvo B.M., Phys. Rep., C41(1978)225;
Boehm F., Vogel P., Physics of Massive Neutrinos: Cambridge Univ. Press, 1987, p.27, p.121;
Bilenky S.M., Petcov S.T., Rev. of Mod. Phys., 1977, v.59, p.631.
2. C. Gonzalez-Garcia 31-st ICHEP, Amsterdam, July 2002.
3. Rosen S.P., Lecture Notes on Mass Matrices, LASL preprint, 1983.
4. Beshtoev Kh.M., JINR Communication D2-2001-292, Dubna, 2001;
5. Beshtoev Kh.M., Internet Publ. hep-ph/0103274.
6. Beshtoev Kh.M., JINR Commun. E2-92-195, Dubna, 1992.
7. Glashow S.L.- Nucl. Phys., 1961, vol.22, p.579 ;
Weinberg S.- Phys. Rev. Lett., 1967, vol.19, p.1264 ;
Salam A.- Proc. of the 8th Nobel Symp., edited by N. Svarthholm (Almgvist and Wiksell, Stockholm) 1968, p.367.
8. Phys. Rev D66, N1, 2002, Review of Particle Physics.